

ON KERNEL THEOREMS FOR (LF)-SPACES

A. G. SMIRNOV

ABSTRACT. A convenient technique for proving kernel theorems for (LF)-spaces (countable inductive limits of Fréchet spaces) is developed. The proposed approach is based on introducing a suitable modification of the functor of the completed inductive topological tensor product. Using such modified tensor products makes it possible to prove kernel theorems without assuming the completeness of the considered (LF)-spaces. The general construction is applied to proving kernel theorems for a class of spaces of entire analytic functions arising in nonlocal quantum field theory.

1. INTRODUCTION

Let X and Y be sets and F , G , and H be locally convex spaces consisting of functions defined on X , Y , and $X \times Y$ respectively. Suppose the function $(x, y) \rightarrow f(x)g(y)$ belongs to H for any $f \in F$ and $g \in G$. In applications, it is often important to find out whether the following statement holds:

- (K) For any separately continuous bilinear mapping ψ from $F \times G$ to a Hausdorff complete space $\tilde{H}$, there is a unique continuous linear mapping $\tilde{\psi}: H \rightarrow \tilde{H}$ such that $\psi = \tilde{\psi} \circ \Phi$, where the bilinear mapping $\Phi: F \times G \rightarrow H$ is defined by the relation $\Phi(f, g)(x, y) = f(x)g(y)$.

Statements of this type and their analogues for multilinear mappings are known as kernel theorems. In this paper, we propose a convenient technique for proving results of this type in the case, where all considered spaces are (LF)-spaces (i.e., countable inductive limits of Fréchet spaces).

In the language of topological tensor products [2], (K) means that H can be identified with a dense subspace of the completion $F \hat{\otimes}_i G$ of $F \otimes G$ with respect to the inductive¹ topology. In particular, to prove (K), it suffices to show that H can be identified with $F \hat{\otimes}_i G$. However, this requires proving the completeness of H , which may present difficulty for concrete functional (LF)-spaces (and is actually unnecessary, as we shall see). To circumvent this problem, we introduce a notion of the semi-completed tensor product of (LF)-spaces (see Definition 2). The semi-completed tensor product $F \tilde{\otimes} G$ of (LF)-spaces F and G is an (LF)-space which is canonically identified with a dense subspace of $F \hat{\otimes}_i G$, and, therefore, we can prove (K) by identifying H with $F \tilde{\otimes} G$. Moreover, semi-completed tensor products possess a natural associativity property which turns out to be very useful for the treatment of multilinear mappings.

The research was supported by the grants RFBR 05-01-01049, INTAS 03-51-6346, LSS-4401.2006.2, and the grant MK-1315.2006.1 of the President of Russian Federation.

¹Recall that the inductive (projective) topology on $F \otimes G$ is the strongest locally convex topology on $F \otimes G$ such that the canonical bilinear mapping $(f, g) \rightarrow f \otimes g$ is separately continuous (resp., continuous).

The paper is organized as follows. In Sec. 2, we give the definition of semi-completed tensor products and describe their basic properties. In Sec. 3, we find simple conditions ensuring the coincidence of $F \hat{\otimes} G$ with a given space H in the case, where F , G , and H are functional (LF)-spaces. In Sec. 4, we apply the obtained results to proving kernel theorems for some spaces of entire analytic functions arising in nonlocal quantum field theory (see [6]).

In Secs. 2 and 3, we assume that the considered vector spaces are either real or complex. The ground field ($\mathbb{R}$ or $\mathbb{C}$) is denoted by $\mathbb{K}$.

2. SEMI-COMPLETED TENSOR PRODUCTS OF (LF)-SPACES

A Hausdorff locally convex space F is called an (LF)-space if there exist a sequence (F_i) of Fréchet spaces and a sequence p_i of continuous mappings from F_i to F such that $F = \bigcup_i \text{Im } p_i$ and the topology of F coincides with the inductive topology with respect to the mappings p_i . The following important result was proved by Grothendieck (see Théorème A of [2]).

Lemma 1. *Let F be a Hausdorff locally convex space, (F_i) be a sequence of Fréchet spaces, and (p_i) be a sequence of continuous mappings from F_i to F . Let $\tilde{F}$ be a Fréchet space and $p: \tilde{F} \rightarrow F$ be a continuous mapping such that $\text{Im } p \subset \bigcup_i \text{Im } p_i$. Then there is an index i such that $\text{Im } p \subset \text{Im } p_i$ and if p_i is injective, then there is a continuous mapping $\tilde{p}: \tilde{F} \rightarrow F_i$ such that $p = p_i \circ \tilde{p}$.*

It follows from Lemma 1 that the topology of an (LF)-space F coincides with the inductive topology with respect to an arbitrary countable family of continuous mappings of Fréchet spaces into F provided that the images of these mappings cover F .

Let F be a Hausdorff locally convex space and $\tilde{F}$ be a Fréchet space. We say that $\tilde{F}$ is a Fréchet subspace of F (notation $\tilde{F} \prec F$) if $\tilde{F} \subset F$ and the inclusion mapping $\tilde{F} \rightarrow F$ is continuous. The relation $\prec$ is a partial order on the set $\mathfrak{F}_F$ of all Fréchet subspaces of F . In fact, $\mathfrak{F}_F$ is a lattice: the least upper bound $F_1 \vee F_2$ of $F_1, F_2 \in \mathfrak{F}_F$ is the space $F_1 + F_2$ endowed with the inductive topology with respect to the inclusion mappings $F_{1,2} \rightarrow F_1 + F_2$; the greatest lower bound $F_1 \wedge F_2$ is the space $F_1 \cap F_2$ endowed with the projective topology with respect to the inclusion mappings $F_1 \cap F_2 \rightarrow F_{1,2}$. In particular, the set $\mathfrak{F}_F$ is directed. By Lemma 1, a countable set $U \subset \mathfrak{F}_F$ is cofinal in $\mathfrak{F}_F$ if and only if $F = \bigcup_{\tilde{F} \in U} \tilde{F}$. If F is an (LF)-space, then $\mathfrak{F}_F$ contains a countable cofinal subset. Indeed, let a sequence (F_i) of Fréchet spaces and a sequence (p_i) of continuous linear mappings from F_i to F be such that $F = \bigcup_i \text{Im } p_i$. Let $\tilde{F}_i$ be the space $\text{Im } p_i$ endowed with the inductive topology with respect to the mapping p_i . Then $\tilde{F}_i \in \mathfrak{F}_F$ for all i and, therefore, $\tilde{F}_i$ form the required subset.

For a locally convex space F , we denote by $\hat{F}$ the Hausdorff completion of F . Given locally convex spaces $F_1, \dots, F_n$, the tensor product $F_1 \otimes \dots \otimes F_n$ endowed by the inductive (resp., projective) topology will be denoted by $F_1 \otimes_i \dots \otimes_i F_n$ (resp., by $F_1 \otimes_\pi \dots \otimes_\pi F_n$). The Hausdorff completions of $F_1 \otimes_i \dots \otimes_i F_n$ and $F_1 \otimes_\pi \dots \otimes_\pi F_n$ will be denoted by $F_1 \hat{\otimes}_i \dots \hat{\otimes}_i F_n$ and $F_1 \hat{\otimes}_\pi \dots \hat{\otimes}_\pi F_n$ respectively. If $F_1, \dots, F_n$ are Fréchet spaces, then the inductive and projective topologies on $F_1 \otimes \dots \otimes F_n$ coincide. In this case, we shall omit the indices i and π in the notation for tensor products.

Definition 2. Let $F_1, \dots, F_n$ be (LF)-spaces. The semi-completed tensor product $F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n$ of $F_1, \dots, F_n$ is defined to be the Hausdorff space associated with the inductive limit

$$\lim_{(\tilde{F}_1, \dots, \tilde{F}_n) \in \mathfrak{F}_{F_1} \times \dots \times \mathfrak{F}_{F_n}} \tilde{F}_1 \hat{\otimes} \dots \hat{\otimes} \tilde{F}_n.$$

Each of the sets $\mathfrak{F}_{F_1}, \dots, \mathfrak{F}_{F_n}$ contains a countable cofinal subset. It hence follows that $\mathfrak{F}_{F_1} \times \dots \times \mathfrak{F}_{F_n}$ also contains a countable cofinal subset and, therefore, $F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n$ is an (LF)-space. For $\tilde{F} = (\tilde{F}_1, \dots, \tilde{F}_n) \in \mathfrak{F}_{F_1} \times \dots \times \mathfrak{F}_{F_n}$, let $\iota_{\tilde{F}}$ be the natural continuous linear mapping from $\tilde{F}_1 \otimes \dots \otimes \tilde{F}_n$ to $F_1 \otimes \dots \otimes F_n$ determined by the inclusion mappings $\tilde{F}_i \rightarrow F_i$, $i = 1, \dots, n$, and $\rho_{\tilde{F}}$ be the canonical continuous linear mapping from $\tilde{F}_1 \hat{\otimes} \dots \hat{\otimes} \tilde{F}_n$ to $F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n$.

Lemma 3. Let $F_1, \dots, F_n$ be (LF)-spaces. There is a unique continuous linear mapping $j: F_1 \otimes \dots \otimes F_n \rightarrow F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n$ such that

$$j\iota_{\tilde{F}} = \rho_{\tilde{F}}\lambda_{\tilde{F}},$$

for any $\tilde{F} = (\tilde{F}_1, \dots, \tilde{F}_n) \in \mathfrak{F}_{F_1} \times \dots \times \mathfrak{F}_{F_n}$, where $\lambda_{\tilde{F}}$ is the canonical mapping from $\tilde{F}_1 \otimes \dots \otimes \tilde{F}_n$ to $\tilde{F}_1 \hat{\otimes} \dots \hat{\otimes} \tilde{F}_n$. The mapping j is a topological isomorphism of $F_1 \otimes \dots \otimes F_n$ onto a dense subspace of $F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n$. There is a unique continuous linear mapping $k: F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n \rightarrow F_1 \hat{\otimes} \dots \hat{\otimes} F_n$ such that $k \circ j$ is the canonical mapping $F_1 \otimes \dots \otimes F_n \rightarrow F_1 \hat{\otimes} \dots \hat{\otimes} F_n$. The mapping k is a topological isomorphism of $F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n$ onto a dense subspace of $F_1 \hat{\otimes} \dots \hat{\otimes} F_n$. For every $f \in F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n$, there is a bounded subset B of $F_1 \otimes \dots \otimes F_n$ such that $f \in \overline{j(B)}$, where the bar means closure.

The mapping j described in Lemma 3 will be called the canonical mapping from $F_1 \otimes \dots \otimes F_n$ to $F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n$. For $f_1 \in F_1, \dots, f_n \in F_n$, we set

$$f_1 \tilde{\otimes} \dots \tilde{\otimes} f_n = j(f_1 \otimes \dots \otimes f_n).$$

The n -linear mapping $(f_1, \dots, f_n) \rightarrow f_1 \tilde{\otimes} \dots \tilde{\otimes} f_n$ will be called the canonical n -linear mapping from $F_1 \times \dots \times F_n$ to $F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n$.

Definition 4. Let $F_1, \dots, F_n$ be (LF)-spaces, H be a locally convex space, and φ be an n -linear mapping from $F_1 \times \dots \times F_n$ to H . We say that φ is (F)-continuous if for any $\tilde{F}_1 \in \mathfrak{F}_{F_1}, \dots, \tilde{F}_n \in \mathfrak{F}_{F_n}$, there are a Fréchet space $\tilde{H}$, a continuous n -linear mapping $\psi: \tilde{F}_1 \times \dots \times \tilde{F}_n \rightarrow \tilde{H}$, and a continuous linear mapping $l: \tilde{H} \rightarrow H$ such that $l \circ \psi$ coincides with the restriction of φ to $\tilde{F}_1 \times \dots \times \tilde{F}_n$.

Obviously, every (F)-continuous n -linear mapping is separately continuous. If H is Hausdorff and complete, then every separately continuous n -linear mapping $\varphi: F_1 \times \dots \times F_n \rightarrow H$ is (F)-continuous. Indeed, for any $\tilde{F}_1 \in \mathfrak{F}_{F_1}, \dots, \tilde{F}_n \in \mathfrak{F}_{F_n}$, the restriction of φ to $\tilde{F}_1 \times \dots \times \tilde{F}_n$ can be decomposed as $l \circ \psi$, where ψ is the canonical n -linear mapping from $\tilde{F}_1 \times \dots \times \tilde{F}_n$ to $\tilde{F}_1 \hat{\otimes} \dots \hat{\otimes} \tilde{F}_n$ and $l: \tilde{F}_1 \hat{\otimes} \dots \hat{\otimes} \tilde{F}_n \rightarrow H$ is a continuous linear mapping.

It easily follows from the above definitions that the canonical n -linear mapping from $F_1 \times \dots \times F_n$ to $F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n$ is (F)-continuous. Moreover, the space $F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n$ has the following universal property:

Lemma 5. Let $F_1, \dots, F_n$ be (LF)-spaces, H be a Hausdorff locally convex space, and φ be an (F)-continuous n -linear mapping from $F_1 \times \dots \times F_n$ to H . Then there

is a unique continuous linear mapping $l: F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n \rightarrow H$ such that

$$\varphi(f_1, \dots, f_n) = l(f_1 \tilde{\otimes} \dots \tilde{\otimes} f_n), \quad f_1 \in F_1, \dots, f_n \in F_n.$$

Theorem 6. *Let $F_1, \dots, F_n$ be (LF) -spaces. For every $1 \leq m < n$, there is a unique topological isomorphism $F_1 \tilde{\otimes} \dots \tilde{\otimes} F_n \simeq (F_1 \tilde{\otimes} \dots \tilde{\otimes} F_m) \tilde{\otimes} (F_{m+1} \tilde{\otimes} \dots \tilde{\otimes} F_n)$ taking $f_1 \tilde{\otimes} \dots \tilde{\otimes} f_n$ to $(f_1 \tilde{\otimes} \dots \tilde{\otimes} f_m) \tilde{\otimes} (f_{m+1} \tilde{\otimes} \dots \tilde{\otimes} f_n)$ for any $f_1 \in F_1, \dots, f_n \in F_n$.*

3. TENSOR PRODUCTS OF FUNCTIONAL (LF) -SPACES

Given a locally convex space F , we denote by F' and $\langle \cdot, \cdot \rangle$ the continuous dual of F and the canonical bilinear form on $F' \times F$ respectively. We denote by F'_σ , F'_τ , and F'_b the space F' endowed with its weak topology, Mackey topology, and strong topology respectively.

Let X and Y be sets and F and G be locally convex spaces consisting of functions defined on X and Y respectively. We denote by $\Pi(F, G)$ the linear space consisting of all functions $h(x, y)$ on $X \times Y$ such that $h(x, \cdot) \in G$ for every $x \in X$ and the function $h_v(x) = \langle v, h(x, \cdot) \rangle$ belongs to F for every $v \in G'$.

Lemma 7. *Let X and Y be sets and F , G , and H be Hausdorff complete locally convex spaces consisting of scalar functions defined on X , Y , and $X \times Y$ respectively. Let F be B -complete, G be nuclear, and the topologies of F , G , and H be stronger than that of simple convergence. Suppose the function $(x, y) \rightarrow f(x)g(y)$ on $X \times Y$ belongs to H for every $f \in F$ and $g \in G$ and the bilinear mapping $\Phi: F \times G \rightarrow H$ taking (f, g) to this function is continuous. Then Φ induces an injective continuous linear mapping $F \hat{\otimes}_\pi G \rightarrow H$ whose image coincides with $\Pi(F, G)$.*

Proof. Without loss of generality, we can assume that $H = \mathbb{K}^{X \times Y}$. Let $\Phi_*: F \hat{\otimes}_\pi G \rightarrow H$ be the continuous linear mapping determined by Φ . As usual, let $\mathfrak{B}_e(F'_\sigma, G'_\sigma)$ denote the space of separately continuous bilinear forms on $F'_\sigma \times G'_\sigma$ equipped with the biequicontinuous convergence topology (i.e., the topology of the uniform convergence on the sets of the form $A \times B$, where A and B are equicontinuous sets in F' and G' respectively). Let S be the natural continuous linear mapping $F \hat{\otimes}_\pi G \rightarrow \mathfrak{B}_e(F'_\sigma, G'_\sigma)$ which takes $f \otimes g$ to the bilinear form $(u, v) \rightarrow \langle u, f \rangle \langle v, g \rangle$. Since G is nuclear, S is a topological isomorphism (see [2], Chapitre 2, Théorème 6 or [3]). Further, let T be the linear mapping $\mathfrak{B}_e(F'_\sigma, G'_\sigma) \rightarrow \mathbb{K}^{X \times Y}$ defined by the relation $(Tb)(x, y) = b(\delta_x, \delta_y)$, $b \in \mathfrak{B}_e(F'_\sigma, G'_\sigma)$ (if $x \in X$ and $y \in Y$, then δ_x and δ_y are the linear functionals on F and G such that $\langle \delta_x, f \rangle = f(x)$ and $\langle \delta_y, g \rangle = g(y)$; they are continuous because the topologies of F and G are stronger than the topology of simple convergence). Obviously, T is continuous and TS coincides with Φ_* on $F \otimes G$. By continuity, we have $\Phi_* = TS$ everywhere on $F \hat{\otimes}_\pi G$. Moreover, T is injective because δ -functionals are weakly dense in F' and G' . To prove the statement, we therefore have to show that $\text{Im } T = \Pi(F, G)$.

Let $b \in \mathfrak{B}_e(F'_\sigma, G'_\sigma)$ and $h = Tb$. The bilinear form b determines two linear mappings $L_1: F' \rightarrow G$ and $L_2: G' \rightarrow F$ such that $b(u, v) = \langle v, L_1 u \rangle = \langle u, L_2 v \rangle$ for any $u \in F'$ and $v \in G'$. For $x \in X$ and $y \in Y$, we have $h(x, y) = \langle \delta_y, L_1 \delta_x \rangle = (L_1 \delta_x)(y)$, i.e., $h(x, \cdot) = L_1 \delta_x$. Further, for $v \in G'$ and $x \in X$, we have $h_v(x) = \langle v, h(x, \cdot) \rangle = \langle v, L_1 \delta_x \rangle = \langle \delta_x, L_2 v \rangle = (L_2 v)(x)$. Hence $h_v = L_2 v$ belongs to G and, therefore, $h \in \Pi(F, G)$. Thus, we have the inclusion $\text{Im } T \subset \Pi(F, G)$.

We now prove the converse inclusion. Let $h \in \Pi(F, G)$ and $L: G'_\tau \rightarrow F$ be the linear mapping taking $v \in G'$ to h_v . We claim that the graph $\mathcal{G}$ of L is closed.

It suffices to show that if an element of the form $(0, f)$ belongs to the closure $\bar{\mathcal{G}}$ of $\mathcal{G}$, then $f = 0$. Suppose the contrary that there is $f_0 \in F$ such that $f_0 \neq 0$ and $(0, f_0) \in \bar{\mathcal{G}}$. Let $x_0 \in X$ be such that $f_0(x_0) \neq 0$ and let the neighborhood U of f_0 be defined by the relation $U = \{f \in F : |\langle \delta_{x_0}, f - f_0 \rangle| < |f_0(x_0)|/2\}$. Let $V = \{v \in G' : |\langle v, h(x_0, \cdot) \rangle| < |f_0(x_0)|/2\}$. If $f \in U$ and $v \in V$, then we have $|h_v(x_0)| < |f_0(x_0)|/2 < |f(x_0)|$. Hence the neighborhood $V \times U$ of $(0, f_0)$ does not intersect $\mathcal{G}$. This contradicts to the assumption that $(0, f_0) \in \bar{\mathcal{G}}$, and our claim is proved. Being nuclear and complete, G is semireflexive and hence G'_τ is barrelled. We can therefore apply the closed graph theorem ([3], Theorem IV.8.5) and conclude that L is continuous. Let the bilinear form b on $F' \times G'$ be defined by the relation $b(u, v) = \langle u, h_v \rangle$. The continuity of L implies that $b \in \mathfrak{B}_e(F'_\sigma, G'_\sigma)$. Since $b(\delta_x, \delta_y) = h(x, y)$, we have $h = Tb$. Thus, $h \in \text{Im } T$ and the lemma is proved. $\square$

Theorem 8. *Let X and Y be sets, F , G , and H be (LF)-spaces consisting of scalar functions on X , Y , and $X \times Y$ respectively. Suppose the topologies of F , G , and H are stronger than that of simple convergence and G can be covered by a countable family of its nuclear Fréchet subspaces. Let the following conditions be satisfied:*

- (i) *For every $f \in F$ and $g \in G$, the function $(x, y) \rightarrow f(x)g(y)$ on $X \times Y$ belongs to H and the bilinear mapping $\Phi: F \times G \rightarrow H$ taking (f, g) to this function is (F) -continuous.*
- (ii) *For any $h \in H$, one can find Fréchet subspaces $\tilde{F} \subset F$ and $\tilde{G} \subset G$ such that $h(x, \cdot) \in \tilde{G}$ for every $x \in X$ and the function $h_v(x) = \langle v, h(x, \cdot) \rangle$ belongs to $\tilde{F}$ for every $v \in \tilde{G}'$.*

Then Φ induces the topological isomorphism $F \tilde{\otimes} G \simeq H$.

Proof. Let $\Phi_*: F \tilde{\otimes} G \rightarrow H$ be the continuous linear mapping determined by Φ . For $\tilde{F} \in \mathfrak{F}_F$ and $\tilde{G} \in \mathfrak{F}_G$, let $\Phi_*^{\tilde{F}, \tilde{G}}$ be the continuous linear mapping $\tilde{F} \tilde{\otimes} \tilde{G} \rightarrow H$ determined by the restriction of Φ to $\tilde{F} \times \tilde{G}$. We have

$$(1) \quad \Phi_*^{\tilde{F}, \tilde{G}} = \Phi_* \rho_{\tilde{F}, \tilde{G}},$$

where $\rho_{\tilde{F}, \tilde{G}}$ is the canonical mapping from $\tilde{F} \hat{\otimes} \tilde{G}$ to $F \tilde{\otimes} G$. Let $\mathcal{N}$ be the subset of $\mathfrak{F}_G$ consisting of all nuclear Fréchet subspaces of G . By the assumption, $\mathcal{N}$ is cofinal in $\mathfrak{F}_G$. Let $h \in H$. Condition (ii) means that $h \in \Pi(\tilde{F}, \tilde{G})$ for some $\tilde{F} \in \mathfrak{F}_F$ and $\tilde{G} \in \mathfrak{F}_G$. Let $\tilde{G}_1 \in \mathcal{N}$ be such that $\tilde{G} \prec \tilde{G}_1$. Then we have $\Pi(\tilde{F}, \tilde{G}) \subset \Pi(\tilde{F}, \tilde{G}_1)$ and, therefore, $h \in \Pi(\tilde{F}, \tilde{G}_1)$. By Lemma 7, we have $\Pi(\tilde{F}, \tilde{G}_1) = \text{Im } \Phi_*^{\tilde{F}, \tilde{G}_1}$ and in view of (1) we conclude that $h \in \text{Im } \Phi_*$. Thus, Φ_* is surjective. By Lemma 7, the mappings $\Phi_*^{\tilde{F}, \tilde{G}}$ are injective for any $\tilde{F} \in \mathfrak{F}_F$ and $\tilde{G} \in \mathcal{N}$. Since $\mathcal{N}$ is cofinal in $\mathfrak{F}_G$, it follows from (1) that the restriction of Φ_* to the image of $\rho_{\tilde{F}, \tilde{G}}$ is injective for any $\tilde{F} \in \mathfrak{F}_F$ and $\tilde{G} \in \mathfrak{F}_G$. This implies the injectivity of Φ_* because the spaces $\text{Im } \rho_{\tilde{F}, \tilde{G}}$ cover $F \tilde{\otimes} G$. We have thus proved that Φ_* is a one-to-one mapping from $F \tilde{\otimes} G$ onto H . Since both $F \tilde{\otimes} G$ and H are (LF)-spaces, we can apply the open mapping theorem ([2], Théorème B) and conclude that Φ_* is a topological isomorphism. The theorem is proved. $\square$

4. APPLICATIONS TO SPACES OF ANALYTIC FUNCTIONS

Let U be a cone in $\mathbb{R}^k$. We say that a cone W is a conic neighborhood of U if W has an open projection² and contains U .

Definition 9. Let $\beta(s)$ be continuous monotone indefinitely increasing convex function on the semi-axis $s \geq 0$ and U be a nonempty cone in $\mathbb{R}^k$. For $B > 0$, the Fréchet space $\mathcal{E}^{\beta,B}(U)$ consists of entire analytic functions on $\mathbb{C}^k$ having the finite norms

$$\|f\|_{U,N,B'} = \sup_{z=x+iy \in \mathbb{C}^k} |f(z)|(1+|x|)^N \exp(-\beta(B'|y|) - \beta(B'\delta_U(x)))$$

for any $B' > B$ and any nonnegative integer N , where $\delta_U(x) = \inf_{x' \in U} |x - x'|$ is the distance from x to U . The space $\mathcal{E}^\beta(U)$ is defined by the relation $\mathcal{E}^\beta(U) = \bigcup_{B>0, W \supset U} \mathcal{E}^{\beta,B}(W)$, where W runs over all conic neighborhoods of U and the union is endowed with the inductive limit topology.

The spaces $\mathcal{E}^\beta(\mathbb{R}^k)$ proved to be useful for the analysis of infinite series in the Wick powers of free fields converging to nonlocal fields [6]. If $\beta(s) = s^{1/(\nu-1)}$, $\nu < 1$, then $\mathcal{E}^\beta(\mathbb{R}^k)$ coincides with the Gelfand-Shilov space $S^\nu(\mathbb{R}^k)$ (see [1] for the definition and properties of Gelfand-Shilov spaces; to avoid confusion, we use the notation S^ν instead of the standard S^β). The spaces $\mathcal{E}^\beta(U)$ over cones are introduced to describe the localization properties of analytic functionals belonging to $\mathcal{E}'^\beta(\mathbb{R}^k)$. More precisely, a closed cone $K \subset \mathbb{R}^k$ is called a carrier cone of $u \in \mathcal{E}'^\beta(\mathbb{R}^k)$ if u has a continuous extension to $\mathcal{E}'^\beta(K)$. The notion of carrier cone replaces the notion of support of a generalized function for elements of $\mathcal{E}'^\beta(\mathbb{R}^k)$. In particular, every $u \in \mathcal{E}'^\beta(\mathbb{R}^k)$ has a uniquely determined minimal carrier cone (this was proved in [7] for the case of the space S^0 ; the general case can be treated in the same way using the estimates for plurisubharmonic functions obtained in [5]).

Here, we shall prove a kernel theorem for the spaces $\mathcal{E}^\beta(U)$. For this, we introduce, in addition to $\mathcal{E}^\beta(U)$, similar spaces associated with finite families of cones.

Definition 10. Let $U_1, \dots, U_n$ be nonempty cones in $\mathbb{R}^{k_1}, \dots, \mathbb{R}^{k_n}$ respectively. We define the space $\mathcal{E}^\beta(U_1, \dots, U_n)$ by the relation

$$\mathcal{E}^\beta(U_1, \dots, U_n) = \bigcup_{B>0, W_1 \supset U_1, \dots, W_n \supset U_n} \mathcal{E}^{\beta,B}(W_1 \times \dots \times W_n),$$

where the union is taken over all conic neighborhoods $W_1, \dots, W_n$ of $U_1, \dots, U_n$ and is endowed with the inductive limit topology.

In what follows, we assume for definiteness that the norm $|\cdot|$ on $\mathbb{R}^k$ is uniform: $|x| = \max_{1 \leq j \leq k} |x_j|$. For any cone U , the space $\mathcal{E}^{\beta,B}(U)$ belongs to the class of the spaces $\mathcal{H}(M)$ introduced in [4]. The space $\mathcal{H}(M)$ is defined³ by a family $M = \{M_\gamma\}_{\gamma \in \Gamma}$ of strictly positive continuous functions on $\mathbb{C}^k$ and consists of all entire analytic functions on $\mathbb{C}^k$ with the finite norms

$$\sup_{z \in \mathbb{C}^k} M_\gamma(z) |f(z)|.$$

²The projection $\text{Pr } W$ of a cone $W \subset \mathbb{R}^k$ is by definition the intersection of W with the unit sphere in $\mathbb{R}^k$; the projection of W is meant to be open in the topology of this sphere.

³The definition of $\mathcal{H}(M)$ given here is slightly less general than that in [4], but it is sufficient for our purposes.

It is supposed that (a) for every $\gamma_1, \gamma_2 \in \Gamma$, one can find $\gamma \in \Gamma$ and $C > 0$ such that $M_\gamma \geq C(M_{\gamma_1} + M_{\gamma_2})$, and (b) there is a countable set $\Gamma' \subset \Gamma$ with the property that for every $\gamma \in \Gamma$, one can find $\gamma' \in \Gamma'$ and $C > 0$ such that $CM_\gamma \leq M_{\gamma'}$ (a family of functions on $\mathbb{C}^k$ satisfying (a) and (b) will be called a defining family of functions). Given a cone $U \subset \mathbb{R}^k$ and $B > 0$, we define the family $M^{U,B}$ of functions on $\mathbb{C}^k$ indexed by the set $\mathbb{Z}_+ \times (B, \infty)$ by the relation

$$(2) \quad M_{N,B'}^{U,B}(x+iy) = (1+|x|)^N \exp(-\beta(B'|y|) - \beta(B'\delta_U(x))), \quad N \in \mathbb{Z}_+, B' > B.$$

Then all above conditions are satisfied and we have

$$(3) \quad \mathcal{H}(M^{U,B}) = \mathcal{E}^{\beta,B}(U).$$

By Lemma 12 of [4], $\mathcal{H}(M)$ is a nuclear Fréchet space if the following conditions are satisfied:

- (I) For every $\gamma \in \Gamma$, there is $\gamma' \in \Gamma$ such that $M_\gamma(z)/M_{\gamma'}(z)$ is integrable on $\mathbb{C}^k$ and tends to zero as $|z| \rightarrow \infty$.
- (II) For every $\gamma \in \Gamma$, there are $\gamma' \in \Gamma$, a neighborhood of the origin $\mathcal{B}$ in $\mathbb{C}^k$, and $C > 0$ such that $M_\gamma(z) \leq CM_{\gamma'}(z + \zeta)$ for any $z \in \mathbb{C}^k$ and $\zeta \in \mathcal{B}$.

It is straightforward to verify that the family $M^{U,B}$ satisfies (I) and (II). The space $\mathcal{E}^{\beta,B}(U)$ is therefore nuclear for any $B > 0$ and cone $U \subset \mathbb{R}^k$.

Let $M = \{M_\gamma\}_{\gamma \in \Gamma}$ and $N = \{N_\omega\}_{\omega \in \Omega}$ be defining families of functions on $\mathbb{C}^{k_1}$ and $\mathbb{C}^{k_2}$ respectively. We denote by $M \otimes N$ the family formed by the functions

$$(M \otimes N)_{\gamma\omega}(z_1, z_2) = M_\gamma(z_1)N_\omega(z_2), \quad (\gamma, \omega) \in \Gamma \times \Omega.$$

Clearly, if $M \otimes N$ is a defining family of functions. The following result was proved in [4].

Lemma 11. *Let $M = \{M_\gamma\}_{\gamma \in \Gamma}$ and $N = \{N_\omega\}_{\omega \in \Omega}$ be defining families of functions on $\mathbb{C}^{k_1}$ and $\mathbb{C}^{k_2}$ respectively and let $h \in \mathcal{H}(M \otimes N)$. Suppose N satisfies conditions (I) and (II). Then $h(z, \cdot) \in \mathcal{H}(N)$ for every $z \in \mathbb{C}^{k_1}$ and the function $h_v(z) = \langle v, h(z, \cdot) \rangle$ belongs to $\mathcal{H}(M)$ for all $v \in \mathcal{H}'(N)$.*

Let $V_1 \subset \mathbb{R}^{k_1}$ and $V_2 \subset \mathbb{R}^{k_2}$ be nonempty cones. It follows from (2) and the monotonicity of β that

$$M_{N,2B'}^{V_1 \times V_2, 2B}(z_1, z_2) \leq M_{N,B'}^{V_1,B}(z_1) M_{N,B'}^{V_2,B}(z_2) \leq M_{2N,B'}^{V_1 \times V_2, B}(z_1, z_2), \quad z_{1,2} \in \mathbb{C}^{k_{1,2}},$$

for any $B' > B > 0$ and $N \in \mathbb{Z}_+$. We hence have continuous inclusions

$$(4) \quad \mathcal{E}^{\beta,B}(V_1 \times V_2) \subset \mathcal{H}(M^{V_1,B} \otimes M^{V_2,B}) \subset \mathcal{E}^{\beta,2B}(V_1 \times V_2).$$

Lemma 12. *Let $U_1, \dots, U_n$ be nonempty cones in $\mathbb{R}^{k_1}, \dots, \mathbb{R}^{k_n}$ respectively. Let $1 \leq m < n$ and $\Phi: \mathcal{E}^\beta(U_1, \dots, U_m) \times \mathcal{E}^\beta(U_{m+1}, \dots, U_n) \rightarrow \mathcal{E}^\beta(U_1, \dots, U_n)$ be the bilinear mapping defined by the relation*

$$\Phi(f, g)(z_1, \dots, z_n) = f(z_1, \dots, z_m) g(z_{m+1}, \dots, z_n).$$

Then Φ is (F)-continuous and induces the topological isomorphism

$$\mathcal{E}^\beta(U_1, \dots, U_m) \tilde{\otimes} \mathcal{E}^\beta(U_{m+1}, \dots, U_n) \simeq \mathcal{E}^\beta(U_1, \dots, U_n).$$

Proof. We check that the spaces $F = \mathcal{E}^\beta(U_1, \dots, U_m)$, $G = \mathcal{E}^\beta(U_{m+1}, \dots, U_n)$, and $H = \mathcal{E}^\beta(U_1, \dots, U_n)$ and the bilinear mapping Φ satisfy conditions (i) and (ii) of Theorem 8. Let $B > 0$ and $W_1, \dots, W_n$ be conic neighborhoods of $U_1, \dots, U_n$ respectively. Let $V_1 = W_1 \times \dots \times W_m$ and $V_2 = W_{m+1} \times \dots \times W_n$. In view

of (3) it follows from (4) that Φ induces a continuous bilinear mapping $\mathcal{E}^{\beta,B}(V_1) \times \mathcal{E}^{\beta,B}(V_2) \rightarrow \mathcal{E}^{\beta,B}(V_1 \times V_2)$. This implies that Φ is (F)-continuous and (i) is fulfilled. Let $h \in \mathcal{E}^{\beta,B}(W_1 \times \dots \times W_n)$. It follows from (4) and Lemma 11 that (ii) will be satisfied if we set $\tilde{F} = \mathcal{E}^{\beta,B}(V_1)$ and $\tilde{G} = \mathcal{E}^{\beta,B}(V_2)$. The lemma is proved. $\square$

The next result follows from Lemma 12 and Theorem 6 by induction on n .

Theorem 13. *Let $U_1, \dots, U_n$ be nonempty cones in $\mathbb{R}^{k_1}, \dots, \mathbb{R}^{k_n}$ respectively and $\Phi: \mathcal{E}^{\beta}(U_1) \times \dots \times \mathcal{E}^{\beta}(U_n) \rightarrow \mathcal{E}^{\beta}(U_1, \dots, U_n)$ be the bilinear mapping defined by the relation*

$$\Phi(f_1, \dots, f_n)(z_1, \dots, z_n) = f_1(z_1) \dots f_n(z_n).$$

Then Φ is (F)-continuous and induces the topological isomorphism

$$\mathcal{E}^{\beta}(U_1) \tilde{\otimes} \dots \tilde{\otimes} \mathcal{E}^{\beta}(U_n) \simeq \mathcal{E}^{\beta}(U_1, \dots, U_n).$$

REFERENCES

- [1] I. M. Gelfand and G. E. Shilov, *Generalized functions*, Vol. 2, Academic Press, New York–London, 1967.
- [2] A. Grothendieck, *Produits tensoriels topologiques et espaces nucléaires*, Mem. Amer. Math. Soc. **16** (1955).
- [3] H. Schaefer, *Topological Vector Spaces*, Springer, Berlin–Heidelberg–New York, 1981.
- [4] A. G. Smirnov, “On topological tensor products of functional Fréchet and DF spaces,” arXiv:math.FA/0512611.
- [5] A. G. Smirnov, “On localization properties of highly singular generalized functions,” arXiv:math-ph/0512062, accepted for publication in Theor. Math. Phys.
- [6] A. G. Smirnov, M. A. Soloviev, *Wick power series converging to nonlocal fields*, Theor. Math. Phys. **127** (2001) 632.
- [7] M. A. Soloviev, “Carrier cones of analytic functionals,” arXiv:math-ph/0507011.

I. E. TAMM THEORY DEPARTMENT, P. N. LEBEDEV PHYSICAL INSTITUTE, LENINSKY PROSPECT
53, MOSCOW 119991, RUSSIA
E-mail address: smirnov@lpi.ru